

1 Chapter 1

2 Introduction

3 Fisheries Science is the study of exploited fish stocks [3, 14]. With the advent of vessel monitoring
4 systems, new assessment methods are needed that can make use of higher resolution catch and
5 effort data. Some of these systems are capable of recording a time stamp and global positioning
6 system coordinates for each unit of effort. Looking at continuously collected data has challenges,
7 the analyst has a choice to bin the data by weeks, month, years, or even hours or minutes. Depletion
8 estimation of abundance is an established method, [7, 19], and seasonal catch and effort data are
9 becoming easier to obtain. Modern computing tools make fitting dynamic models much easier; it
10 is easy to imagine the use of depletion models on daily, weekly, or monthly catch and effort series.
11 One good example of a fishery where this idea has been applied is the Moroccan octopus fishery
12 [22], where recruitment and harvest were estimated on a weekly time step. For instance, weekly
13 models have seen some use in invertebrate fisheries [9, 11, 22]. Given the number of weeks in a year,
14 significantly more data are available in systems with weekly collected data compared to systems
15 which collect data annually. Despite this, relatively few tools have been developed that use weekly
16 data to study the processes of recruitment, mortality, harvest, and growth. Some well studied
17 invertebrate fisheries that do use weekly or monthly models include the Northern Prawn Fishery in
18 Australia [10], and neon flying squid in the Pacific [6].

19 Such weekly depletion models, particularly when coupled with size composition data, are able
20 to provide estimates of realized, as opposed to targeted, exploitation rates. With periodic size (or
21 age composition) data, natural mortality and seasonal patterns in vulnerability can also be studied.

22 Most fisheries have some combination of input and output regulations imposed on them. How-
23 ever, there are few tools that are capable of integrating input and output regulations, market forces,
24 physical limitations (e.g., weather), population dynamics, and the growth in size of individuals and
25 the life cycles of harvested species into an understanding of temporal patterns in the exploitation
26 rate. This is largely due to few within season assessment models having been developed[22].
27 Few management strategy evaluation models are capable of evaluating commonly-used manage-
28 ment choices, such as seasonal closures, size restrictions, and restrictions on the number of licenses.
29 Within season models would allow for the simulation of non-regulatory constraints and drivers of
30 fishermen behavior. Seasonal models may be particularly important to some of the valuable
31 effort-controlled fisheries, such as the Dungeness crab fishery and the Atlantic lobster fishery.

32 Management strategy evaluation has been increasing in popularity as a tool to evaluate the
33 performance of fishery regulations [21]. However, it is rare that management strategy evaluations
34 are used on within season management. Expanding the management strategy evaluation concept to
35 include a more complete picture of the different limits on exploitation rates is needed to understand
36 how management choices are likely to affect the social, economic, and ecological dimensions of
37 fisheries. There is increasing understanding that fishery management is about "people, not fish" [12,
38 25]; this can only be especially important in fisheries with complex seasonal patterns in the amount
39 of latent (capable of being deployed, but still tied up on shore) effort [25]. Within season depletion
40 models allow for the simulated evaluation of seasonal management regimes that are impossible under
41 annual operating models.

42 This dissertation develops a within season depletion model for the Dungeness crab fisheries. It
43 then uses the model to simulate different input controls including changes to the number of vessels
44 in the fleet and rules regarding opening and closing the fishery to evaluate how much catch is lost
45 due to softshell handling.

46 1.1 Dungeness Crab Fishery

47 For this analysis the Dungeness Crab Fishery in the Hecate Strait was used as a case study, however
48 Dungeness crab fisheries are conducted from Alaska to California [8]. This section provides the
49 relevant background information on the biology, economics, and history of the fishery.

50 1.2 Biology

51 Dungeness crab (*Cancer magister* [1] or *Metacarcinus magister* [2]) have a complicated life history
52 including a several month period as pelagic zooplankton [23]. While this thesis is not the place for
53 a detailed exposition of everything known about their life history, it is important to review some
54 relevant details that are important to management. There has been some effort made mapping how
55 zoea are currently distributed along the pacific coast [15, 16]. In the late spring or early summer,
56 zoea moult into the megalope life stage and the currents bring them close to shore [18]. During
57 their first two years, they moult often, but as the crabs grow in size they moult less frequently until
58 reaching three years of age, when they only moult about once a year [24]. In subsequent chapters,
59 crab growth is discussed in greater detail.

60 To reproduce, a male must be wider than a female crab, while females can only receive a sperm
61 plug when their shell is soft and can fertilize multiple (annual) batches of eggs from a single sperm
62 plug [20] [13]. Since the Dungeness crab fishery exclusively targets large males, concerns have
63 been raised that recruitment may be impacted by females failing to find large males to mate with.
64 However, Hankin et al. [13] did not find any evidence for that.

65 1.3 Coastwide history of the Dungeness crab fishery.

66 Whole books could be written about the development of the Dungeness crab fishery, detailing the
67 different types of traps used by decade, the bottom trawl fishery, the transition from selling canned
68 crab meat to live crab, and countless other shifts in climate, ecology, technology, and economics that
69 shaped this fishery. This thesis does not attempt to undertake such an endeavor. In particular there
70 are some well-studied cycles in California [4, 5, 16] and long-term time series of annual landings in
71 British Columbia that are important context to the analyses undertaken in subsequent chapters.

72 1.3.1 California Cycles

73 Dungeness crab landings in Northern California have historically fluctuated in a nearly cyclic man-
74 ner, with six or seven years of high production followed by about four years of low production. A
75 summary of these cycles is shown in table 1.1. The California cycles were the subject of significant
76 study because they occurred with such regularity. It is of note that there was almost an order of
77 magnitude difference between the years of high production and the years of low production. Sev-

78 eral hypotheses have been proposed to explain the periods of high and low productivity including
 79 oceanographic conditions, lags in fishing effort, and cannibalism [4], [5], [16]. Looking at long-term
 80 records Johnson et al. [17] shows that somewhat regular periods of high and low production occurred
 in Washington and Oregon as well.

Years	Average landings (lbs/year)
1955-1961	9,755,310
1961-1965	1,922,148
1965-1971	10,474,386
1971-1975	1,281,444
1975-1982	13,330,590
1982-1984	4,587,442

Table 1.1: Crab landing Cycles in California

81

82 1.3.2 Historical Catches in British Columbia

83 These figures come from data in the book written by Jim Boutillier at Fisheries and Oceans Canada
 84 (D.F.O); I do not know if the book was ever properly published. It is not entirely clear how to map
 85 the data to geography as maps showing what area in text corresponds to what area in reality are
 86 difficult to find. That said, Figure 1.1 corresponds to the area near the Skeena river which is near
 87 the Hecate Strait.

88 1.4 Management

89 Dungeness crab are managed in British Columbia under a collection of regulations: only male crabs
 90 larger than 165 mm with hard shells can be harvested [8] and there are restrictions on the number
 91 of vessels that can fish in a given area, the size of the vessels, and the number of traps per vessel
 92 [26]. Finally, to avoid intensive fishing during time periods when many soft shell males are present,
 93 there are area dependent rules to close the fishery during a moult.

94 There are a few things to consider about the management regime: the size restriction is effectively
 95 a sex restriction, and few female crabs are larger than 165mm. Also the major markets for B.C.
 96 crab are China and the United States of America, so any discussion about bringing soft shell crabs
 97 to market would have to consider the logistics of live shipment of soft shell crab.




Figure 1.1: Effort as predicted by catch per trap haul.

98 1.5 Economics

99 Yonis [26] points out that there were 9 vessels larger than 52 meters that could deploy up to 1200
 100 traps in the Hecate Strait (known officially as Area A). There were also 10 vessels between 42.5
 101 meters and 52 meters that could deploy between 800 and 1000 traps. The remaining vessels were
 102 less than 42.5 meters and could deploy up to 600 traps.




Figure 1.2: Landings in B.C.

103 Yonis [26] estimated that the fleet average crew share was \$97,835 with the “bottom tier” earning
 104 \$29,097 and the “top tier” earning \$198,072. This is much higher than the crew shares in the other
 105 areas, which had a mean of \$24,063.

106 The crab fleet is one of the least diversified fleets in British Columbia, 80% of the fleet has only
 107 a single license and 96% of the fleet has less than two different fishing licenses [26]. This is primarily
 108 due to the fact that crab boats are specifically built for crabbing.




Figure 1.3: Landings in B.C.

109 The top third of the fleet in Area A accounted for almost 60% of the catch in 2007[26]. The
 110 bottom third of the fleet accounted for only 13% of the catch.

111 Yonis [26] points out that price fluctuates over the course of a year. In spring of 2008 (March,
 112 April, and May) the price per kilogram was about \$10 but in July it dropped down to about \$8.
 113 From personal communication I have discovered that there are two hypothesis to explain why prices
 114 seem to drop when Area A opens. The first hypothesis is that Area A lands so many crab that the

115 spike in supply drops the price of crab. The second hypothesis is that shipping live crab from the
116 Hecate Strait results in an inferior product commanding a lower price.

117 The United States of America, China, and Hong Kong are the leading markets for crab Yonis
118 [26]. There may be tacit agreements to time the Washington and Oregon crab fisheries out of sync
119 with the B.C. crab fishery to maintain market prices, although it is unclear how important this is
120 compared to weather and the seasonal dynamics of hard shell crab.

121 The following chapters examine the Dungeness crab fishery in the Hecate Strait in British
122 Columbia as a case study. There are four chapters in this thesis: a chapter that completes an
123 empirical analysis of effort patterns over the course of the fishing season, a chapter that develops
124 and fits a weekly depletion model to the weekly catch and effort data, a chapter that extends the
125 depletion model to incorporate size composition data, and a chapter that simulates three different
126 rules involving the opening and closing dates of the fishery. Incorporating size composition data
127 allows for the investigation of natural mortality and time changing patterns in vulnerability. Vessel
128 Monitoring System data opens up the potential to study fine scale temporal, and in some cases
129 spatial, relationships between fishing effort, resource abundance, and catch; new tools are needed to
130 make the most of this relatively modern source of data. This thesis seeks to develop some of these
131 tools.

References

- 133 [1] (2019). Integrated taxonomic information system (itis). <http://www.itis.gov/>. → pages 3
- 134 [2] (2019). World register of marine species (worms). =<http://www.marinespecies.org>. Accessed:
135 2019-10-25. → pages 3
- 136 [3] Beverton, R. J. and Holt, S. J. (1993). On the Dynamics of Exploited Fish Populations. The
137 Blackburn press. → pages 1
- 138 [4] Botsford, L. W. and Hobbs, R. C. (1995). Recent advances in the understanding of cyclic
139 behavior of dungeness crab (cancer magister) populations. In ICES Marine Science Symposia,
140 volume 199, pages 157–166. Copenhagen, Denmark: International Council for the Exploration of
141 the Sea, 1991. → pages 3, 4
- 142 [5] Botsford, L. W., Methot Jr, R. D., and Johnston, W. E. (1983). Effort dynamics of the northern
143 california dungeness crab (cancer magister) fishery. Canadian Journal of Fisheries and Aquatic
144 Sciences, 40(3):337–346. → pages 3, 4
- 145 [6] Chen, X., Chen, Y., Tian, S., Liu, B., and Qian, W. (2008). An assessment of the west win-
146 ter–spring cohort of neon flying squid (*ommastrephes bartramii*) in the northwest pacific ocean.
147 Fisheries Research, 92(2–3):221 – 230. → pages 1
- 148 [7] DeLury, D. B. (1947). On the estimation of biological populations. Biometrics, 3(4):145–167.
149 → pages 1
- 150 [8] DFO (2010). Pacific Region Integrated Fisheries Management Plan. Technical report, Fisheries
151 and Oceans Canada. → pages 2, 4
- 152 [9] Dichmont, C., Deng, A., Punt, A., Ellis, N., Venables, W., Kompas, T., Ye, Y., Zhou, S., and
153 Bishop, J. (2008). Beyond biological performance measures in management strategy evaluation:
154 Bringing in economics and the effects of trawling on the benthos. Fisheries Research, 94(3):238 –
155 250. Advances in the Analysis and Application of Harvest Policies in the Management of Fisheries.
156 → pages 1
- 157 [10] Dichmont, C., Punt, A., Deng, A., Dell, Q., and Venables, W. (2003). Application of a weekly
158 delay-difference model to commercial catch and effort data for tiger prawns in australia’s northern
159 prawn fishery. Fisheries Research, 65(1):335 – 350. Life Histories, Assessment and Management
160 of Crustacean Fisheries. → pages 1
- 161 [11] Dichmont, C. M., Deng, A. R., Punt, A. E., Venables, W., and Haddon, M. (2006). Management
162 strategies for short-lived species: The case of australia’s northern prawn fishery: 1. accounting for
163 multiple species, spatial structure and implementation uncertainty when evaluating risk. Fisheries
164 research, 82(1-3):204–220. → pages 1

- 165 [12] Grafton, Q. R., Hilborn, R., Squires, D., and Williams, M. J. (2010). Marine Fisheries
166 Conservation and Management, chapter 1. Oxford University Press. → pages 2
- 167 [13] Hankin, D. G., Butler, T. H., Wild, P. W., and Xue, Q. L. (1997). Does intense fishing
168 on males impair mating success of female dungeness crabs. Canadian Journal of Fisheries and
169 Aquatic Sciences, 54(3):655–669. → pages 3
- 170 [14] Hilborn, R. and Walters, C. J. (1992). Quantitative Fisheries Stock Assessment Choice Dynamics
171 and Uncertainty. Kluwer Academic Publishers. → pages 1
- 172 [15] Jamieson, G. S. and Phillips, A. (1993). Megalopal Spatial Distribution and Stock Separation in
173 Dungeness Crab (*Cancer magister*). Canadian Journal of Fisheries and Aquatic Science, 50:416–
174 429. → pages 3
- 175 [16] Johnson, D. F., Botsford, L. W., Methot Jr, R. D., and Wainwright, T. C. (1986a). Wind
176 stress and cycles in dungeness crab (*cancer magister*) catch off california, oregon, and washington.
177 Canadian Journal of Fisheries and Aquatic Sciences, 43(4):838–845. → pages 3, 4
- 178 [17] Johnson, D. F., Botsford, L. W., Methot Jr., R. D., and Wainwright, T. C. (1986b). Wind
179 stress and cycles in dungeness crab (*cancer magister*) catch off california, oregon, and washington.
180 Canadian Journal of Fisheries and Aquatic Sciences, 43(4):838–845. → pages 4
- 181 [18] Johnson, J. and Shanks, A. L. (2002). Time series of the abundance of the post-larvae of the
182 crabscancer *magister* and *cancer* spp. on the southern oregon coast and their cross-shelf transport.
183 Estuaries, 25(6):1138–1142. → pages 3
- 184 [19] Leslie, P. H. and Davis, D. H. S. (1939). An attempt to determine the absolute number of rats
185 on a given area. Journal of Animal Ecology, 8(1):94–113. → pages 1
- 186 [20] Oh, S. J. and Hankin, D. G. (2004). The Sperm Plug Is a Reliable Indicator of Mating Success
187 in Female Dungeness Crabs, *Cancer Magister*. Journal of Crustacean Biology, 24(2):314–326. →
188 pages 3
- 189 [21] Punt, A. E., Butterworth, D. S., de Moor, C. L., De Oliveira, J. A. A., and Haddon, M. (2016).
190 Management strategy evaluation: best practices. Fish and Fisheries, 17(2):303–334. → pages 2
- 191 [22] Robert, M., Faraj, A., McAllister, M. K., and Rivot, E. (2010). Bayesian State-Space modelling
192 of the De Lury depletion model; strengths and limitations of the method, and application to the
193 Moroccan octopus fishery. ICES Journal of Marine Science, 67:1272–1290. → pages 1, 2
- 194 [23] Shanks, A., Roegner, C. C., and Miller, J. (2010). Using Megalopae Abundance To Predict
195 Future Commercial Catches of Dungeness Crabs (*Cancer Magister*) in Oregon. Technical report,
196 CalCOFI 51. → pages 3
- 197 [24] Wainwright, T. C. and Armstrong, D. A. (1993). Growth patterns in the dungeness crab
198 (*cancer magister dana*): Synthesis of data and comparison of models. Journal of Crustacean
199 Biology, 13(1):36–50. → pages 3
- 200 [25] Walters, C. J. and Martell, J. (2004). Fisheries Ecology and Management. Princeton University
201 Press. → pages 2
- 202 [26] Yonis, R. (2010). The Economics of British Columbia’s Crab Fishery: Socio-economic Profile,
203 Viability, and Market Trends. Technical report, Fisheries and Oceans Canada. → pages 4, 5, 6,
204 7, 8